

Dingo

Canis familiaris dingo



The dingo is a primitive canid related to wolves and coyote. The dingo was not a part of the ancestral fauna of Australia. Though its origins are not clear, it is thought to have arrived in Australia 3500–4000 years ago.

It is the largest mammalian carnivore remaining in mainland Australia, and as such fills an important ecological niche. Females weigh about 12 kg and males 15 kg.

The dingo has been regarded as a serious predator of domestic stock since early European settlement in Australia.

Since European settlement domestic dogs have been released or escaped into the environment to cross with dingoes. These hybrids or crosses are colloquially called wild dogs (*Canis lupus familiaris*). Often the term wild dog covers both dingoes and dingo hybrids.

Wild dogs predate on livestock, native fauna and domestic pets.



Legal requirements

The dingo is a category 3, 4, 5 and 6 restricted invasive animal under the *Biosecurity Act 2014*. It must not be moved, kept, fed, given away, sold, or released into the environment. A permit to deal with dingoes maybe given in limited circumstances.

The Act requires everyone to take all reasonable and practical measures to minimise the biosecurity risks associated with invasive animals under their control. This is called a general biosecurity obligation (GBO).

At a local level, each local government must have a biosecurity plan that covers invasive animals in its area. This plan may include actions to be taken on certain species. Some of these actions may be required under local laws. Contact your local government for more information.

Description

Red, ginger and sandy-yellow are the dominant coat colours, though dingoes can also be pure white, black and tan or solid black.

It is not difficult to distinguish between most dingoes and hybrids. The presence of domestic genes is suggested by broken colours—brindling and patchiness in the normally pure white feet and chest patch and sable colouration (black hairs along the back and sides).

Dingoes have a more heavily boned skull and larger teeth (especially the canine) than domestic dogs of similar size.

Life cycle

Dingoes have only one breeding season per year (usually April to June), whereas domestic bitches have two or more oestrus cycles per year. However, unless seasons are particularly favourable, or human sources of food are intentionally or inadvertently provided, feral domestic dogs are unlikely to successfully rear two litters per year.

After a nine-week gestation, dingo pups (usually four to six) are born in a hollow log or cave den. Bitches tend to use the same den each year. Pups are suckled at four to six weeks and generally weaned at four months. When large enough to travel, pups are taken from the den to kills, and other dens may be used. The range of pups is increased as they are moved from den to den. In this way the pups are gradually moved around the bitch's home range.

Independence may occur as early as six months of age when parents abandon them, but this results in high juvenile mortality. Pups that become independent around 12 months appear to disperse voluntarily. Being larger and more experienced, mortality is then usually low.

Where dingoes live alone or in small groups (most pastoral and semi-settled areas), mature females will breed successfully each year.

By contrast, dominant female infanticide results in only one litter being successfully raised each year within groups containing several adult females (e.g. undisturbed areas such as the Simpson Desert). The dominant (alpha)

female will kill all pups of the other females, and then use subordinate females to suckle and rear her litter.

Methods of spread

Dingoes in an undisturbed area generally belong to discrete packs (3–12 members), which occupy long-term, non-overlapping territories. The group rarely moves as a pack—rather, members meet and separate again throughout the day. Dingoes are most gregarious during the breeding season.

There is overlap of home ranges within a group. In contrast, boundaries between groups are more rigid, actively defended and infrequently crossed.

Olfactory communication (smell) is important in dingo social organisation. Dingo droppings are deposited along pads in specific areas where other dingoes will encounter them (creek crossings, intersections of roads and fences).

These 'scent posts' appear to delineate the home range boundary and act as a warning to neighbouring groups and individuals.

This strong site attachment of dingoes is contrary to the notion commonly held by property owners that dingoes will travel large distances to kill stock.

Habitat and distribution

Dingo numbers are believed to be higher today than in pre-European times. This is thought to be due to increased food availability via the introduced rabbit and cattle carcasses, and the development of permanent waters in arid areas of the state. Dingoes/wild dogs are present in all parts of the state.

The distribution of the wild dog in relation to purebred dingoes varies throughout the state. In far western areas, most dingoes sighted appear to be 'pure', with characteristic white points and broad heads. Closer to settled areas a greater number of feral domestic dogs produce a generally hybrid population. It has been estimated that dingoes are 50% pure in south-eastern Queensland and 90–95% pure in south-western and central Queensland.

Radio tracking studies show dingoes occupy a discrete area known as a 'home range'. The dingo visits the edge of this area frequently.

The home range can vary in size according to the productivity of the country—from 9 km² in rainforest areas to 300 km² on the Nullarbor Plain.

The edge of the home range is commonly associated with a major topographic feature (e.g. an escarpment, a major ridge or stream).

The home range is not used uniformly. Activity is centred on areas with highest food density.

Hunting movement is slow and exploratory, in contrast to frequent rapid movement around the home range boundary.

Pads follow well defined paths and are most likely associated with sociality and home range boundary maintenance. Activity is highest at dusk and dawn.

Diet

Dietary research of stomach content and faecal scats has shown dingoes are opportunistic predators.

Medium-size animals such as kangaroos, wallabies, rabbits and possums consistently form the major part of the dingo diet.

Studies by the Western Australia Agriculture Protection Board show dingoes in undisturbed refuge areas killed and ate kangaroos strictly according to need.

On grazing country, however, 'dingoes harassed, bit or killed sheep in large numbers, often without eating any'. The consumption of these sheep carcasses was the exception rather than the rule. Even kangaroos in these areas were sometimes killed in 'play' type behaviour rather than for food.

Such dietary studies could suggest dingo predation of domestic stock is low. There is, however, a need for caution in using such studies to assess dingo impact on stock.

Grouping increases foraging efficiency and appears necessary to exploit larger prey. Dingoes cooperating in groups are more successful in hunting kangaroos than lone dingoes are. While lone dingoes can easily kill sheep, it is less likely a solitary dingo would successfully attack a calf in the presence of a defending cow.

Disease threat

Dingoes are vectors of canid diseases (e.g. distemper, parvovirus) and parasites. The hydatid parasite *Echinococcus granulosus* is a major problem of dogs and domestic stock. It can cause illness and occasionally death in humans.

The dingo could pose a serious risk if the exotic disease rabies was introduced to Australia.

Beneficial considerations

The establishment of watering points during post-European settlement has resulted in a huge increase in the kangaroo population, with consequent strong pasture competition with domestic livestock.

Though it is widely accepted that sheep production is near impossible in the presence of dingoes, many cattle producers will tolerate dingoes because of their believed suppression of kangaroo numbers.

Research has shown that in some cases the dingo has the potential to mitigate population growth of native species during abundant seasons and it could also be an important limiting factor for many feral animal populations (e.g. feral pigs and goats).

There is some evidence that destruction of the dingo could cause increases in other pests to the grazing industry and result in widespread degradation of environmentally sensitive areas. However, this has not been proven.

Further information

Further information is available from your local government office, or by contacting Biosecurity Queensland on 13 25 23 or visit biosecurity.qld.gov.au.





This fact sheet is developed with funding support from the Land Protection Fund.

Fact sheets are available from Department of Agriculture and Fisheries (DAF) service centres and our Customer Service Centre (telephone 13 25 23). Check our website at biosecurity.qld.gov.au to ensure you have the latest version of this fact sheet. The control methods referred to in this fact sheet should be used in accordance with the restrictions (federal and state legislation, and local government laws) directly or indirectly related to each control method. These restrictions may prevent the use of one or more of the methods referred to, depending on individual circumstances. While every care is taken to ensure the accuracy of this information, DAF does not invite reliance upon it, nor accept responsibility for any loss or damage caused by actions based on it.