



Red-eared slider turtle

Trachemys scripta (subspecies *elegans*)

Steve Csurhes and Clare Hankamer

First published 2012

Updated 2016



Queensland
Government

© State of Queensland, 2016.

The Queensland Government supports and encourages the dissemination and exchange of its information. The copyright in this publication is licensed under a Creative Commons Attribution 3.0 Australia (CC BY) licence.



You must keep intact the copyright notice and attribute the State of Queensland as the source of the publication.

Note: Some content in this publication may have different licence terms as indicated.

For more information on this licence visit <http://creativecommons.org/licenses/by/3.0/au/deed.en>

Contents

Summary	4
Introduction	5
Identity and taxonomy	5
Description	5
Biology and ecology	7
Reproduction and dispersal	7
Origin and distribution	8
Status in Australia	9
Status in Queensland	10
Preferred habitat	11
History as a pest elsewhere	11
Uses	12
Pest potential in Queensland	12
Pest potential of related taxa	15
Bibliography	16

Summary

The red-eared slider turtle is a small freshwater turtle native to the Mississippi Valley in the southern United States.

Tens of millions of red-eared sliders have been sold worldwide as pets. As a result, the species has naturalised in numerous countries. While perhaps tending to become most abundant in heavily modified waterways, there is considerable evidence that red-eared sliders can negatively affect locally native turtle species. The International Union for Conservation of Nature (IUCN) has listed the red-eared slider among 100 examples of the ‘world’s worst invasive species’. Pet turtles can also transmit salmonella to people, especially children.

Despite strong restrictions on possession and sale in Queensland, small naturalised populations of red-eared sliders have been detected in coastal south-eastern Queensland. These populations have been reduced to very low levels via a concerted eradication campaign over the last eight years, but it is difficult to claim complete eradication. The continued dumping of unwanted pets means that surveillance and control will be ongoing.

Based on the evidence presented in this risk assessment, it seems reasonable to suggest red-eared sliders are a ‘high-risk’ species in Queensland. The species has a range of attributes considered to confer significant pest potential, including:

- a history as a major pest overseas
- climatic suitability to much of Queensland
- a broad geographic range
- a generalist diet
- high fecundity.

This study suggests there is sufficient evidence to conclude red-eared sliders pose a significant threat to native freshwater turtle species and human health, and that current restrictions on possession and sale are justified.

Introduction

Identity and taxonomy

Species identity: *Trachemys scripta*, subspecies *elegans*

Synonyms: *Chrysemys scripta elegans*, *Chrysemys scripta*, *Emys elegans*, *Emys holbrooki*, *Emys sanguinolenta*, *Pseudemys scripta*, *Pseudemys scripta elegans*, *Testudo scripta*, *Trachemys lineata*

Common names: Red-eared slider turtle

Family: Emydidae

Currently, the red-eared slider turtle is considered a subspecies of *Trachemys scripta* ('common slider' or 'slider'). However, its taxonomy is unclear. In the past, it has been placed in three different genera—*Chrysemys*, *Pseudemys* and *Trachemys* (Gibbons 1990). While the various genera are subject to ongoing review, *T. scripta* is currently split into three subspecies—*T.s. elegans* (red-eared slider), *T.s. scripta* (yellow-bellied slider) and *T.s. troostii* (Cumberland slider) (Seidel 2002).

Hybrids of red-eared sliders and yellow-bellied sliders have been reported in the southern United States (Somma et al. 2009a), hybrids of red-eared sliders and *T. taylorii* in Mexico (Seidel 2002), and hybrids of red-eared sliders and native terrapins in the Bahamas (Ardastra Gardens, Zoo and Conservation Center 2010). There is potential for hybridisation with other congeners, notably *T. decussata angusta* (Cayman Islands) (Lever 2003; GISD 2009), *Trachemys decorata* and *Trachemys stejnegeri vicina* (Caribbean) (Powell et al. 2000), and *Trachemys gaigeae* (New Mexico) (Seidel et al. 1999; Stuart 2000). Hybridisation is common in closely related freshwater turtle species, and intergeneric and interclade hybrids exist (Parham et al. 2001; Iverson et al. 2007; Brophy et al. 2006).

Description

Red-eared sliders are a medium-sized freshwater turtle, generally 125–200 mm long, but can grow up to 350 mm long (Figure 1). Females are usually larger than males (plastron length 150–195 mm, compared to 90–100 mm for males) (Ernst & Barbour 1972). The carapace and skin is olive to brown with yellow stripes or spots. While specimens in captivity tend to have clean shells, the shells of wild specimens are usually covered by a layer of algae, obscuring their distinctive colours and patterns.



Figure 1. Red-eared slider turtle

Key diagnostic features include:

- a red stripe behind each eye (Figure 1)
- an ability to withdraw its neck straight back into its shell (Australian turtle species withdraw their necks sideways)
- long claws and a dome-shaped shell (native turtles have shorter claws and a flattened shell).

Occasionally, the distinctive red stripe is missing or expressed only as a red patch on the top of the head (Frisby 2010). Older males can also become melanistic and difficult to identify (Zipcode Zoo 2010).

Similar native turtle species include:

- *Chelodina longicollis*—common snake-necked turtle, eastern long-necked turtle or stinker, which emits an offensive smell when disturbed (occurs in eastern Australia from northern Queensland to southern Australia)
- *Chelodina oblonga*—oblong turtle (occurs in south-western Western Australia)
- *Myuchelys latisternum*—saw-shelled turtle (occurs in Cape York to northern New South Wales)
- *Emydura macquarii macquarii*—Murray River turtle (occurs in south-eastern Australia from Nepean River, New South Wales, to Pine Rivers, southern Queensland) (Department of Agriculture and Food 2009).

The pet trade has developed two colour morphs of red-eared sliders—a pastel (lighter colour with varying amounts of red and yellow) and an albino (bright yellow as a juvenile with colour fading as the turtle ages) (Frisby 2010).

The two other subspecies—*T.s. scripta* (yellow-bellied slider) and *T.s. troostii* (Cumberland slider)—are similar to red-eared sliders but have a yellow band behind each eye. The Cumberland slider also has a wider post-orbital stripe that is never entirely one colour; more usually yellow and orange fading to a darker orange-red away from the eye (Wikipedia 2010a; Bringsøe 2006). It can appear similar to the yellow-bellied slider, but has fewer wider stripes on its head, neck and legs (Ernst et al. 1994). For more details refer to Bringsøe (2006) and Ernst et al. (1994).

Biology and ecology

Red-eared slider turtles are active during the day (diurnal), resting on the bottom or floating on the surface at night (Department of Agriculture and Food 2009). They can remain submerged for a considerable time.

Adult red-eared sliders tend to be herbivores, whereas juveniles tend to be more omnivorous. Both prey on molluscs and invertebrates opportunistically—notably snails, insects and their larvae, worms, shrimp and crayfish. They are known to eat fish, frogs' eggs and tadpoles, and water snakes. They eat a variety of aquatic plants and algae, including a number of noxious aquatic weeds such as water hyacinth (Ernst et al. 1994; Brown et al. 1995; Feldman 2005). While capable of eating small vertebrates—including small reptiles, amphibians, mammals, fish and birds (Scalera 2006)—fish and birds are very rarely part of their diet (Feldman 2005).

Like most aquatic turtles, red-eared sliders have fixed tongues and must eat their food in the water (Wikipedia 2010b).

Reproduction and dispersal

Sexual maturity is reached at 2–5 years of age (Dundee & Rossman 1989). Maximum life-span is generally in the order of 20–30 years, although only 1% of hatchlings reach this age (Feldman 2005). Some references claim maximum age can be up to 75 years (Dundee & Rossman 1989; Department of Food and Agriculture 2009).

Red-eared sliders can wander up to 9 km from water to find suitable habitat, search for a mate or lay eggs (Department of Agriculture and Food 2009). Nests are dug well above water level, usually within 500 m of water, but sometimes up to 1.6 km away (Bringsøe 2006; Department of Agriculture and Food 2009). Soil moisture is important for nesting and maintenance of egg moisture content. Under laboratory conditions, vermiculite with a water potential of -1500 kpa (0.09 g water: 1 g vermiculite) was the driest condition at which the majority of eggs hatched successfully (Tucker & Paukstis 2000).

Females can produce viable eggs for up to 5 years after mating (O'Keefe 2005), although fertility drops to 20% or less after the first year (Feldman 2005). Females can lay up to 3 clutches (approximately 75 eggs) per season, compared to a maximum of 30 eggs (typically 10–15 eggs) per season for Australian native turtles (Cann 1998; Goode & Russell 1968; O'Keefe 2005; Zipcode Zoo 2010). However, there is evidence that up to 6 clutches per year of 2–30 eggs each can be produced at times (Scalera 2006). Eggs are white, 23–44 mm long and weigh 6–15 g (Bringsøe 2006; Dundee & Rossman 1989).

Red-eared sliders exhibit temperature-dependent sex determination, similar to other turtle species. A temperature of 22.5 °C is the lowest constant incubation temperature needed to produce viable hatchlings. However, most emerge deformed or neurologically impaired at this temperature (Ewert & Nelson 1991). Constant incubation temperature must be greater than 28.3 °C to produce females (Cadi et al. 2004), and to produce all females the temperature must be greater than 30.6 °C for 4 hours per day during the middle (third) phase of development (Feldman 2005).

Time to hatching is approximately 68–70 days. Newborn hatchlings are 20–35 mm long (Dundee & Rossman 1989). Hatchling mortality rates are high. Predators include crows, rats and snakes.

Origin and distribution

Red-eared sliders are native to the Mississippi Valley in the southern United States, including Illinois, Missouri, Kansas, Oklahoma, Arkansas, Tennessee, Mississippi, Louisiana, Texas, Kentucky, Ohio and West Virginia (Department of Agriculture and Food 2009; Somma et al. 2009a).

Naturalised populations exist in every continent except Antarctica, including Asia, Australia, Austria, Bahamas, Bahrain, Belgium, Belize, Bermuda, Brazil, British Virgin Islands, Cambodia, Canada, Cayman Islands, Chile, China, Colombia, Costa Rica, Czech Republic, Cyprus, Denmark, Dominican Republic, El Salvador, Estonia, Europe, Faroe Islands, Finland, France, French Polynesia, French West Indies (Grande-Terre, Basse-Terre), Gambia, Germany, Gibraltar, Greece, Greenland, Guadeloupe, Guam, Guatemala, Guyana, Hong Kong, Honduras, Hungary, Iceland, Indonesia, Israel, Italy, Japan, Korea, Democratic People's Republic of Korea, Republic of Latvia, Lithuania, Malaysia, Martinique, Mexico, Micronesia, Federated States of Micronesia, New Caledonia, New Zealand, Nicaragua, Northern Mariana Islands, Norway, Panama, Philippines, Poland, Puerto Rico, Reunion, Singapore, South Africa, Spain, Sri Lanka, Sweden, Switzerland, Taiwan, Thailand, Trinidad and Tobago, United Kingdom, United States, Venezuela and Vietnam (GISD 2009; Reptile Database 2010) (Figure 2).



Figure 2. Global distribution of red-eared slider turtle (GISD 2009)

Status in Australia

Red-eared sliders have naturalised in the Australian Capital Territory and New South Wales (Table 1). Individual specimens have been detected in the wild in Victoria and Western Australia (GISD 2009; Massam et al. 2010).

In Victoria, *T.s. elegans* is listed as a ‘controlled pest’ under the *Catchment and Land Protection Act 1994* (Victorian Government 2010). In New South Wales, *T. scripta* is limited to restricted collections under category 2 of the Non-Indigenous Animals Regulation 2006 (New South Wales Government 2010). In the Australian Capital Territory, *T.s. elegans* is notifiable and prohibited in the *Pest Plants and Animals Act 2005* (Australian Capital Territory Government 2009). In Western Australia, *T.s. elegans* and *T.s. scripta* are prohibited and cannot be imported or kept except for scientific or educational purposes by public facilities (Western Australian Government 2004). In the Northern Territory, *T.s. elegans* is a ‘restricted animal’, banned from entry into the territory and banned from release into the wild (Northern Territory Government 2007).

Table 1. Current status of red-eared slider populations in other states

State	Location	Status	Date discovered
New South Wales	Georges River Catchment	Breeding	n/a
New South Wales	Yeramba Lagoon	Breeding	2006
New South Wales	Wolli Creek Reserve	Single animal (gravid female)	2007
New South Wales	Hawkesbury and Nepean rivers	n/a	n/a
Australian Capital Territory	Dams near Murrumbidgee River	Ongoing trapping underway	2003
Victoria	Blackburn Lake Sanctuary, Melbourne	n/a	2007 (unconfirmed)
Victoria	Ruffey Lake Park, Melbourne	n/a	2007 (unconfirmed)
Victoria	Elsternwick Park, Melbourne	n/a	2007 (unconfirmed)
Western Australia	Tomato Lake, Perth	Single animal	2006
Western Australia	Hyde Park, Perth	Single animal	2006

Source: Department of Agriculture and Food (2009)

Status in Queensland

T. scripta, including its subspecies, was banned from possession and sale in Queensland in 1985.

While long suspected to exist in the illegal pet trade, the first naturalised population of red-eared sliders was detected in Queensland in 2004, in a small number of private dams in the Pine Rivers Shire just north of Brisbane. An eradication campaign, combined with a public awareness campaign, resulted in the detection of two other naturalised populations. A small number of specimens being kept as pets were also relinquished to authorities.

Eradication efforts have focused on three sites (Mango Hill, Burpengary and Rosser Park), all in coastal south-eastern Queensland. The total search area is estimated to be approximately 125 ha (Csurhes, unpublished) (Table 2). The species is currently the target of a successful, but ongoing, eradication program that is estimated to have cost at least \$800 000 so far. Current efforts are focused on detecting very small numbers of wild specimens and preventing release/dumping of additional animals.

Table 2. Approximate search area for red-eared slider turtles in Queensland in January 2010 (land and water area suspected to support naturalised populations of red-eared slider turtles)

	Mango Hill	Burpengary	Rosser Park, Ashmore*	Total (ha)
Land area (ha)	42.09	59.34	11.78	113.21
Water area (ha)	5.40	3.66	3.41	12.47
Total (ha)	47.49	63.00	15.19	125.68

*A wild population at Rosser Park is suspected but not confirmed (a small number of eggs have been recorded at the site).

Source: Csurhes (unpublished) from data supplied by S O'Keefe (Biosecurity Queensland) 2010, pers. comm., 5 January.

In 2010, the cost of eradication of remaining residual populations was estimated to be \$270 000 over the following 3 years (Csurhes 2010). The 2010 study considers eradication to be feasible, given the relatively small size of the infested area (126 ha), the marked decline in capture rate (from 140 animals in 2004 to 2 in 2009), and assuming:

- sufficient resources for monitoring, trapping and removal of eggs at current sites
- the development and use of DNA detection technology to remove the last few specimens potentially in dams
- the dumping of additional animals can be prevented by compliance and public awareness (Csurhes 2010).

Preferred habitat

Preferred habitats include a range of still or slow-moving freshwater ponds and lakes (rarely in flowing water). Red-eared sliders readily tolerate artificial ponds and lakes, and often thrive in dams that have been polluted by excessive organic matter. They can tolerate brackish marshes, human-made canals and ponds in city parks. Tolerance of pollution levels including radioactive, thermal, chemical and organic pollution is higher than in other vertebrates (Gibbons 1990). While occasionally seen in salt water, these are animals washed downriver by stormwater.

Preferred climate is perhaps best described as temperate. However, the species is very adaptable and readily tolerates subtropical and tropical areas, generally where temperatures are between 10 °C and 37 °C (Ramsay et al. 2007). In the northern parts of its native range, survival is limited by winter temperature, in particular depth of soil freezing (Packard et al. 1997). In cold climates, hatchlings can overwinter in their nests. Adults can survive severe winters (-10 °C for extended periods) by hibernation—or more correctly brumation, at the bottom of shallow lakes or ponds and less commonly under banks, in hollow tree stumps or under rocks (Wikipedia 2010b; Department of Agriculture and Food 2009). Animals brumate when air temperatures fall below 10 °C or when water temperature falls below 13 °C. They become active when necessary for food or water, or to bask if temperatures increase (Wikipedia 2010b).

Red-eared sliders have been detected in New Zealand but do not appear to be breeding, due to the cold climate (Feldman 2005).

History as a pest elsewhere

The IUCN has listed the red-eared slider among 100 examples of the world's worst invasive species (Lowe et al. 2000). Red-eared sliders have naturalised on every continent except Antarctica and on numerous islands. The Global Invasive Species Database (2009) lists it as a pest in Australia, Bermuda, Brazil, British Virgin Islands, Canada, Cayman Islands, Dominican Republic, France, Germany, Israel, Ruyuku Islands (Japan), Latvia, Poland, Puerto Rico, Singapore, South Africa, Spain, Taiwan and Thailand. In Spain, it is considered one of the most harmful invasive species (GEIB 2006). Despite being native to parts of the United States, it has spread to most states, including Hawaii (Somma et al. 2009b). It was introduced into most countries via the pet trade (GISD 2009; WWF 2010)

Import into New Zealand was banned in 1965. Import into the European Union was banned in 1997 (Bringsøe 2006; Feldman 2005).

There is considerable evidence that red-eared sliders have negative impacts on native turtles and aquatic ecosystems in general. For example, in Singapore, red-eared sliders (now the most commonly observed aquatic turtle in the country) appear to be posing a significant threat to the survival of critically endangered river terrapins (*Batagur baska*) (Blog.nus 2010). Elsewhere, the genetic integrity of rare and threatened *Trachemys* species, such as *Trachemys gaigeae* (Mexican plateau slider) could be at risk, due to hybridisation with red-eared sliders (IUCN 2010; Seidel et al. 1999; Stuart 2000).

Uses

The red-eared slider is the world's most commonly traded live reptile, mainly for use as a pet (Telecky 2001; TRAFFIC 1998; Warwick et al. 1990; WWF 2010). The United States currently exports 8 million red-eared sliders per annum (WWF 2010), with 52 million exported from 1989–1997. Demand was fuelled in the late 1980s by the children's television characters, the 'Teenage Mutant Ninja Turtles' (Somma et al. 2009a). In 1996, red-eared sliders comprised 88% of total trade in reptiles (TRAFFIC 1998). The sheer volume of trade in sliders has facilitated the species' rapid naturalisation across temperate and subtropical countries of the world (Spinks et al. 2003).

Red-eared sliders are popular pets due to low price, small size and easy maintenance. However, they can live for up to 50 years in captivity and, when mature, can inflict painful bites. These traits result in large numbers being dumped into the wild (GISD 2009; Luiselli et al. 1997).

In the Philippines, China and Malaysia, red-eared sliders are farmed for food (Matthews 2008; Somma et al. 2009a; TRAFFIC 2008). Import of turtles into California for their meat was banned in 2010 (USA Today 2010).

In China, slider shells are used in traditional medicine (TRAFFIC 2008).

Millions of red-eared sliders have been released into the wild in Asia for Buddhist 'mercy ceremonies' (Feldman 2005). In Vietnam, they have been released into the wild (along with other animals) during the seventh month of the lunar calendar to pray for good luck.

Pest potential in Queensland

Climate-matching software called 'Climatch' (Bureau of Rural Sciences 2009) was applied to predict areas of Queensland where climate is similar to that experienced across the native and naturalised range of the red-eared slider. Much of southern Queensland and parts of coastal eastern Queensland appear highly suitable (Figure 3).




Figure 3. Area of Australia where climate appears suitable for survival of red-eared slider turtles—red and dark orange indicate areas where climate is highly suitable; light orange and yellow indicate areas where climate is marginally suitable; green and blue indicate areas where climate is considered unsuitable for this species

While the potential impacts of invasive species are generally difficult to predict with precision, information on the impacts of red-eared sliders overseas provides a reasonable insight into the most likely impacts in Queensland. Potential impacts could include the following:

- **Negative impact on native turtle species**

Red-eared sliders have several attributes that seem to confer a competitive advantage over locally native turtle species—they mature at a younger age, are more aggressive, have higher fecundity and have a larger adult body size. A range of studies provide evidence that red-eared sliders can compete successfully with native turtles for food, nesting sites and basking sites (Frank & McCoy 1995; Williams 1999; Salzberg 2000; Cadi & Joly 2003; Polo-Cavia et al. 2010; Lindeman 1999; Spinks et al. 2003; Cann 1998; Goode & Russell 1968; Scalera 2006). For example, Cadi and Joly (2003) found that the endangered European pond turtle (*Emys obicularis galloitalica*) shifted its basking activity to lower quality sites when faced with the more aggressive red-eared slider.

There are 32 species of native freshwater turtles in Australia (Georges & Thomson 2010), including several species classified as rare and threatened in Queensland. If red-eared sliders become abundant across their full potential range in Queensland, they could compete for food resources with locally native turtle species. However, the exact impacts, and the degree to which red-eared sliders might invade less disturbed wetland ecosystems, are difficult to predict.

Hybridisation occurs frequently in turtles, including intrageneric hybridisation (Brophy et al. 2006; Parham et al. 2001; Iverson et al. 2007). Indeed, there has been some debate over new species identifications in the family Geoemydidae and whether they are in fact hybrids produced from turtle farms in China, where multiple species have been housed together (Parham et al. 2001). As yet there is no evidence of interfamilial hybridisation (Brophy et al. 2006) and this seems unlikely. All Australian native turtles belong to the family Chelidae (except *Carettochelys insculpta*, which belongs to the Carettochelyidae). Hence, hybridisation is unlikely to be an issue. It must be noted that *T.s. elegans* and *T.s. scripta* could hybridise in the wild leading to possibly more vigorous genotypes.

- **Negative impact on prey items**

Since red-eared sliders are omnivorous (particularly juveniles), they could have an impact on a range of aquatic prey items, including rare amphibians (O’Keefe 2005; Lever 2003).

- **Negative impact on aquatic ecosystems generally**

Evidence from the United Kingdom suggests red-eared sliders are a threat to nesting waterbirds, taking over nests for basking sites and damaging and preying on eggs and hatchlings (Department of Agriculture and Food 2009). However, the full ecological impact of red-eared sliders is unknown (Ramsay et al. 2007; Bringsøe 2006).

- **Negative impact on human health**

In a controlled study in Connecticut, pet turtles were reported to cause 25% of human salmonellosis cases. Similar results were recorded in another study in New Jersey in 1970–1971 (23% turtle-associated cases, compared to 6% in the control group) (Lamm et al. 1972). Turtles were responsible for an average of 14% of cases per year between 1970 and 1976 (Cohen et al. 1980). Fatalities in children have been reported (Rosenstein et al. 1965; Center for Disease Control 2007). In the state of Washington, a ban on pet turtles enforced in 1968 proved effective in reducing the number of salmonellosis cases, and recommendations were made to enforce the ban across all states (Lamm et al. 1972). To prevent further outbreaks of salmonella in children, there was a United States-wide ban on the sale of turtles with a carapace length less than 4 inches (the size considered small enough to be placed in a child's mouth) (JAMA 2008; Williams 1999). Although this prohibition dramatically reduced the number of cases (Cohen et al. 1980), baby turtles are still available from pet shops and flea markets in the United States. Multi-state outbreaks were reported in 2007–2008, associated with children handling salmonella-infected turtles.

Salmonella can also be transferred to humans when drinking water has been contaminated by turtles (Newberry 1984; Bomford 2008). Use of antibiotics at United States turtle hatcheries has led to more antibiotic resistant strains of salmonella (Ernst et al. 1994).

- **Negative impact on animal health**

Red-eared sliders are often reared in septic conditions. Hence, the risk of spreading diseases and parasites (apart from salmonella) from captive-bred specimens into wild reptile populations is significant (Feldman 2005; Somma et al. 2009a; O'Keefe 2005; Scalera 2006). In the state of Washington, populations of the native turtle *Actinemys marmorata* were seriously impacted by a respiratory infection in 1990 and *T. scripta* was the likely vector (Hays et al. 1999). There is evidence that a malaria-like blood parasite was transferred to two species of native turtles from infected *T.s. elegans* in the Lane Cove River, Sydney (Department of Agriculture and Food 2009).

Based on the evidence presented in this assessment, it seems reasonable to suggest red-eared sliders are a 'high-risk' species in Queensland. The species has a range of attributes considered to confer significant pest potential, including:

- a history as a major pest overseas
- climatic suitability to large parts of Queensland
- a broad geographic range
- a generalist diet
- high fecundity.

Risk assessments in other states also conclude that red-eared sliders are ‘high-risk’ species. Bomford (2008) predicted a ‘serious’ risk of establishment of *T.s. scripta* based on a high climate-match score, evidence of pest status elsewhere and a high establishment score. Similarly, Massam et al. (2010) ranked the species as a ‘high priority’ for management in Australia, to prevent the establishment of additional populations and to eradicate existing populations, and recommended the species be added to a national ‘alert list’.

Pest potential of related taxa

As a consequence of restrictions on sale and possession of red-eared sliders in some countries, closely related species are being sold as alternatives. The yellow-bellied slider (*T.s. scripta*) and the Cumberland slider (*T.s. troostii*) are now replacing the red-eared slider in the European pet trade, and may pose similar threats to the European pond turtle (*Emys orbicularis*) (Bringsøe 2006). Naturalised populations of yellow-bellied sliders have been detected outside the species’ native range in California, Colorado, New York, New Jersey, Pennsylvania, Michigan, Virginia, North Carolina, South Carolina, Wisconsin and perhaps Massachusetts (Somma et al. 2009b), and elsewhere overseas (Ramsay et al. 2007). Hybrids of these subspecies have been recorded in northern Spain (Valdeón et al. 2010).

Bibliography

- AIC 2010, *Illegal trade in flora and fauna and harms to biodiversity*, Australian Institute of Criminology, Canberra, viewed 22 October 2010, <<http://www.aic.gov.au/en/publications/current%20series/rpp/100-120/rpp109/07.aspx>>.
- Alacs, EA & Georges, A 2008, 'Wildlife across our borders: a review of the illegal trade in Australia', *Australian Journal of Forensic Science*, vol. 40, pp. 199–212.
- Ardastra Gardens, Zoo and Conservation Center 2010, Ardastra Gardens, Zoo and Conservation Center, viewed 6 October 2010, <<http://www.ardastra.com/contact.html>>.
- Aresco, MJ 2010, 'Competitive interactions of two species of freshwater turtles, a generalist omnivore and an herbivore, under low resource conditions', *Herpetologica*, vol. 66(3), pp. 259–268.
- Australian Capital Territory Government 2009, *ACT state of the environment 2007*, Office of the Commissioner for Sustainability and the Environment, viewed 3 November 2010, <<http://www.environmentcommissioner.act.gov.au/publications/soe/2007actreport/indicators/pestanimals07>>.
- Australian Wildlife Health Network 2010, *Salmonellosis in reptiles in Australia*, Australian Wildlife Health Network, viewed 12 October 2010, <http://www.wildlifehealth.org.au/AWHN_Admin/ManageWebsite%5CFactSheets%5CUploadedFiles/122/Salmonellosis%20in%20Reptiles%20in%20Australia%205%20Feb%202010%20%281.1%29.pdf>.
- Blog.nus 2010, *The green alien attack!*, Ecological observations in Singapore, viewed 26 October 2010, <<http://blog.nus.edu.sg/lsm2251student/2010/04/14/the-green-alien-attack/>>.
- Bomford, M 2008, *Risk assessment models for establishment of exotic vertebrates in Australia and New Zealand*, Invasive Animals Cooperative Research Centre, Canberra.
- Bouchard, SS & Bjorndal, KA 2005, 'Microbial fermentation in juvenile and adult pond slider turtles, *Trachemys scripta*', *Journal of Herpetology*, vol. 39(2), pp. 321–324.
- Bouchard, SS & Bjorndal, KA 2006, 'Ontogenetic diet shifts and digestive constraints in the omnivorous freshwater turtle *Trachemys scripta*', *Physiology Biochemistry Zoology*, vol. 79, p. 1.
- Bringsøe, H 2006, *Invasive alien species fact sheet—Trachemys scripta*, European Network on Invasive Alien Species, viewed 26 October 2010, <<http://www.nobanis.org>>.
- Brophy, TR, Frair, W & Clark, D 2006, *A review of interspecific hybridization in the order Testudines*, Liberty University, viewed 2 November 2010, <http://digitalcommons.liberty.edu/cgi/viewcontent.cgi?article=1066&context=bio_chem_fac_pubs>.
- Brown, HA, Bury, RB, Darda, DM, Diller, LV, Peterson, CR & Storm, RM 1995, *Reptiles of Washington and Oregon*, Seattle Audubon Society, Seattle.

- Bureau of Rural Sciences 2009, *Climatch*, Department of Agriculture, Fisheries and Forestry, Canberra, <<http://adl.brs.gov.au:8080/Climatch/>>.
- Burgin, S 2007, 'Confirmation of an established population of exotic turtles in urban Sydney', *Australian Zoologist*, vol. 33(3), pp. 379–384.
- Cadi, A, Delmas, V, Prevot-Julliard, A, Joly, P, Pieau, C & Girondot, M 2004, 'Successful reproduction of the introduced slider turtle (*Trachemys scripta elegans*) in the south of France', *Aquatic Conservation: Marine and Freshwater Ecosystems*, vol. 14, pp. 237–246.
- Cadi, A & Joly, P 2003, 'Competition for basking places between the endangered European pond turtle (*Emys orbicularis galloitalica*) and the introduced red-eared slider (*Trachemys scripta elegans*)', *Canadian Journal of Zoology*, vol. 81, pp. 1392–1398.
- Cadi, A & Joly, P 2004, 'Impact of the introduction of the red-eared slider (*Trachemys scripta elegans*) on survival rates of the European pond turtle (*Emys orbicularis*)', *Biodiversity and Conservation*, vol. 13, pp. 2511–2518.
- Cann, J 1998, *Australian freshwater turtles*, Beaumont Publishing, Singapore.
- Center for Disease Control 2007, 'Turtle associated salmonellosis in humans—United States 2006–2007, Center for Disease Control', *MMWR Weekly*, vol. 56(26), pp. 649–652.
- Center for Invasive Species and Ecosystem Health 2010, *Red-eared slider*, Center for Invasive Species and Ecosystem Health, viewed 11 October 2010, <<http://www.invasive.org/species/subject.cfm?sub=14203>>.
- Cohen, ML, Potter, M, Pollard, R & Feldman, RA 1980, 'Turtle associated salmonellosis in the United States; effect of public health action 1970 to 1976', *JAMA*, vol. 243(12), pp. 1247–1249.
- Conant, R & Collins, JT 1991, *Reptiles and amphibians: eastern/central north America*, Houghton Mifflin Company, Boston.
- Cox, MJ, van Dijk, PP, Nabhitabhata, J & Thirakhupt, K 1998, *A photographic guide to snakes and other reptiles of peninsular Malaysia, Singapore and Thailand*, New Holland Publishers (UK) Ltd, London.
- Csurhes, S unpublished, 'Predicting the cost of eradication for Class 1 pest animals in Queensland: small steps towards a useful model', internal report, Biosecurity Queensland, Department of Employment, Economic Development and Innovation, Brisbane.
- Csurhes, S 2010, 'Predicting the cost of eradication for Class 1 pest animals in Queensland', unpublished internal report, Department of Agriculture, Fisheries and Forestry, Brisbane.
- Da Silva E & Blasco M 1995, '*Trachemys scripta elegans* in southwestern Spain', *Herpetological Review*, vol. 26, pp. 133–134.
- Department of Agriculture and Food 2009, *Red-eared slider animal pest alert no. 6/2009*, Department of Agriculture and Food, Western Australia.

Department of Environment, Climate Change and Water 2010, *Seized reptiles moved to Taronga*, media release 16 January, Department of Environment, Climate Change and Water, viewed 28 October 2010, <<http://www.environment.nsw.gov.au/media/DecMedia09011603.htm>>.

Department of Sustainability, Environment, Water, Population and Communities 2009, *Species profile and threats database*, Department of Sustainability, Environment, Water, Population and Communities, viewed 21 October 2010, <http://www.environment.gov.au/cgi-bin/sprat/public/publicthreatenedlist.pl?wanted=fauna#reptiles_vulnerable>.

Dundee, HA & Rossman, DA 1989, *Amphibians and reptiles of Louisiana*, State University Press, Baton Rouge.

Ernst, CH 1990, 'Systematics, taxonomy, variation and geographic distribution of the slider turtle', in CH Ernst, RW Barbour & JE Lovich (eds), *Life history and ecology of the slider turtle*, Smithsonian Institution Press, Washington, pp. 57–67.

Ernst, CH & Barbour, RW 1972, *Turtles of the United States*, University Press of Kentucky, Lexington, Ky.

Ernst, CH & Barbour, RW 1989, *Turtles of the world*, Smithsonian Institution Press, Washington.

Ernst, CH, Lovich, JE & Barbour, RW 1994, *Turtles of the United States and Canada*, Smithsonian Institution Press, Washington and London.

Ewert, MA & Nelson, CE 1991, 'Sex determination in turtles: diverse patterns and some possible adaptive values', *Copeia*, vol. 1, pp. 50–69.

Feldman, ML 2005, 'The red-eared slider turtle (*Trachemys scripta elegans*) in New Zealand', *Proceedings of the 13th Australasian Vertebrate Pest Conference*, Wellington, New Zealand, pp. 96–101.

Frank, JH & McCoy, ED 1995, 'Introduction to insect behavioral ecology: the good, the bad, and the beautiful: non-indigenous species in Florida, invasive adventive insects and other organisms in Florida', *Florida Entomologist*, vol. 78(1), pp. 1–15.

Frisby H 2010, *Red-eared slider: care & feeding of this common pet turtle*, Animal Planet, viewed 26 October 2010, <<http://animal.discovery.com/guides/reptiles/turtles/slider.html>>.

Georges, A & Thomson, S 2010, 'Diversity of Australasian freshwater turtles, with an annotated synonymy and keys to species', *Zootaxa*, vol. 2496, pp. 1–37.

Gibbons, JW (ed.) 1990, *Life history and ecology of the slider turtle*, Smithsonian Institution Press, Washington.

- Global Invasive Species Database (GISD) 2009, *Trachemys scripta elegans*, GISD, Invasive Species Specialist Group, viewed 11 October 2010, <<http://www.issg.org/database/species/ecology.asp?si=71&fr=1&sts=&lang=EN>>.
- Goode, J & Russell, R 1968, 'Incubation of eggs of three species of chelid tortoises, and notes on their embryological development', *Australian Journal of Zoology*, vol. 16, pp. 749–761.
- Grosse, AM, Sterrett, SC & Maerz, JC 2010, 'Effects of turbidity on the foraging success of the eastern painted turtle', *Copeia*, vol. 2010(3), pp. 463–467.
- Grupo Especialista en Invasiones Biológicas (GEIB) 2006, *TOP 20: las 20 especies exóticas invasoras más dañinas presentes en España*, serie técnica no. 2, GEIB.
- Hays, DW, McAllister, KR, Richardson, SA & Swinson DW 1999, *Washington State recovery plan for the western pond turtle*, Washington Department of Fish and Wildlife, Olympia.
- Henderson WR 2009, *Workshop proceedings: risk assessment processes for import and keeping of exotic vertebrates in Australia*, Canberra. Invasive Animals Cooperative Research Centre, Canberra.
- Holman, JA 1994, 'Status of the red-eared slider turtle *Trachemys scripta elegans* (Weid) in Michigan: a preliminary report', *Michigan Academician*, vol. 26, pp. 471–477.
- Integrated Taxonomic Information System (ITIS) 2010, *Trachemys scripta elegans* (*Wied-Neuwied, 1839*), taxonomic serial no. 173823, ITIS, viewed 6 October 2010, <<http://www.itis.gov/servlet/SingleRpt/SingleRpt>>.
- International Union for Conservation of Nature (IUCN) 2010, *IUCN red list of threatened species*, version 2010.3, IUCN, downloaded 21 October 2010, <www.iucnredlist.org>.
- International Union for Conservation of Nature (IUCN) Tortoise and Freshwater Turtle Specialist Group 2007, *Turtles in trouble*, IUCN Tortoise and Freshwater Turtle Specialist Group, viewed 21 October 2010, <<http://www.iucn-tftsg.org/trouble/>>.
- Iverson, JB, Brown, RM, Akre, TS, Near, TJ, Minh, L, Thomson, RC & Starkey, DE 2007, 'In search of the tree of life for turtles, in HB Shaffer, NN FitzSimmons, A Georges & AGJ Rhodin (eds), 'Defining turtle diversity: proceedings of a workshop on genetics, ethics, and taxonomy of freshwater turtles and tortoises', *Chelonian Research Monographs*, vol. 4, pp. 85–106.
- Journal of the American Medical Association (JAMA) 2008, 'Multi-state outbreak of human *Salmonella* infections associated with exposure to turtles—United States 2007–2008', *JAMA*, vol. 299(16), pp. 1892–1894, <<http://jama.ama-assn.org/cgi/reprint/299/16/1892>>.
- Justis C 2010, 'It's winter—where are the turtles?', *The Tennessee Conservationist Magazine*, vol. LXXVI(1) January/February, viewed 28 October 2010, <http://www.state.tn.us/environment/tn_conservationist/turtles.shtml>.

- Lamm, SH, Taylor Jr, A, Gangarosa, EJ, Anderson, HW, Young, W, Clark, MH & Bruce AR 1972, 'Turtle-associated salmonellosis. I. An estimation of the magnitude of the problem in the US, 1970-71', *American Journal Epidemiology*, vol. 95, pp. 511-517.
- Levell, JP 2000, 'Commercial exploitation of Blanding's turtle, *Emydoidea blandingii*, and the wood turtle, *Clemmys insculpta*, for the live animal trade', *Chelonian Conservation and Biology*, vol. 3, pp. 665-674.
- Lever, C 2003, *Naturalized reptiles and amphibians of the world*, Oxford University Press, United Kingdom.
- Lindeman, PV 1999, 'Aggressive interactions during basking among four species of emydid turtles', *Journal of Herpetology*, vol. 33, pp. 214-219.
- Lowe S, Browne M & Boudlejas S 2000, '100 of the world's worst invasive alien species', *Aliens*, vol. 12, pp. 1-12.
- Luiselli, L, Capula M, Capizzi, D, Filippi, E, Jesus, VT & Ainbaldi, C 1997, 'Problems for conservation of pond turtles (*Emys obicularis*) in central Italy: is the introduced red-eared slider (*Trachemys scripta*) a serious threat?', *Chelonian Conservation and Biology*, vol. 2, pp. 417-419.
- Manning, B & Grigg, GC 1997, 'Basking is not of thermoregulatory significance in the 'basking' freshwater turtle *Emydura signata*', *Copeia*, vol. 1997(3), pp. 579-584.
- Maroochy Waterwatch Inc. 2010, *Pest management*, Maroochy Waterwatch Inc., viewed 28 October 2010, <http://www.maroochycatchmentcentre.org.au/catchment/?page_id=75>.
- Martins, TLF, Brooke, MdeL, Hilton, GM, Farnsworth, S, Gould, J & Pain, DJ 2006, 'Costing eradications of alien mammals from islands', *Animal Conservation*, vol. 9, pp. 439-444.
- Massam, M, Kirkpatrick, W & Page, A 2010, *Assessment and prioritisation of risk for forty introduced animal species*, Invasive Animals Cooperative Research Centre, Canberra.
- Matthews, S 2008, *Tropical Asia invaded: the growing danger of invasive alien species*, Global Invasive Species Programme, viewed 26 October 2010, <<http://www.gisp.org/casestudies/showcasestudy.asp?id=212&MyMenuItem=casestudies&worldmap=&country>>.
- Minton Jr., SA 2001, *Amphibians & reptiles of Indiana*, revised 2nd edition, Indiana Academy of Science, Indianapolis.
- Mount, RH 1975, *The reptiles and amphibians of Alabama*, Auburn Printing Co., Auburn.
- Newberry, R 1984, 'The American red-eared terrapin in South Africa', *African Wildlife*, vol. 38, pp. 186-189.
- New South Wales Government 2010, *Non-Indigenous Animals Regulation 2006*, New South Wales Government, viewed 3 November 2010, <<http://www.legislation.nsw.gov.au>>.

Norris, A & Low, T 2005, *Review of the management of feral animals and their impact on biodiversity in the rangelands: a resource to aid NRM planning*, Pest Animal Control Cooperative Research Centre, Canberra.

Northern Territory Government 2007, *Restricted species*, Department of Natural Resources, Environment and the Arts, viewed 3 November 2010, <<http://www.nt.gov.au/nreta/wildlife/permits/restricted.html>>.

Obst, FJ 1983, *Schmuckschildkroten, die Gattung Chrysemys*, A Ziemsen Verlag, Wittenberg Lutherstadt, Germany.

O’Keefe, S 2005, ‘Investing in conjecture: eradicating the red-eared slider in Queensland’, *13th Australasian Vertebrate Pest Conference*, Wellington, New Zealand.

Packard, GC, Tucker, JK, Nicholson, D & Packard, MJ 1997, ‘Cold tolerance in hatching slider turtles (*Trachemys scripta*)’, *Copeia*, vol. 2, pp. 339–345.

Parham, JF, Simison, WB, Kozak, KH, Feldman, CR & Shi, H 2001, ‘New Chinese turtles: endangered or invalid? A reassessment of two species using mitochondrial DNA, allozyme electrophoresis and known-locality specimens’, *Animal Conservation*, vol. 4, pp. 357–367.

Polo-Cavia, N, López, P & Martín, J 2010, ‘Competitive interactions during basking between native and invasive freshwater turtle species’, *Biological Invasions*, vol. 12(7), pp. 2141–2152.

Powell, R, Ottenwalder, JA, Incháustegui, SJ, Henderson, RW & Glor, RE 2000, ‘Amphibians and reptiles of the Dominican Republic: species of special concern’, *Oryx*, vol. 34(2), pp. 118–128.

Ramsay, NF, Ng, PKA, O’Riordan, RM & Chou, LM 2007, ‘The red-eared slider (*Trachemys scripta elegans*) in Asia: a review’, in F Gherardi, *Biological invaders in inland waters: profiles, distribution, and threats*, Springer Publishing, The Netherlands, pp. 161–174.

Reptile Database 2010, *Trachemys scripta*, Reptile Database, viewed 8 October 2010, <<http://reptile-database.reptarium.cz/species.php?genus=Trachemys&species=scripta>>.

Rosenstein, BJ, Russo, P & Hinchliffe, MC 1965, ‘Family outbreak of salmonellosis traced to a pet turtle’, *New England Journal of Medicine*, vol. 272, pp. 960–961.

Salzberg, A 2000, ‘The cage papers. the Norway rat of the turtle world’, *Reptile & Amphibian Hobbyist*, vol. 5(8), p. 84.

Scalera, R 2006, *Trachemys scripta*, Delivering Alien Invasive Species Inventories for Europe, viewed 9 October 2010, <<http://www.europe-aliens.org/speciesFactsheet.do?speciesId=50003#>>.

Seidel, ME 2002, ‘Taxonomic observations on extant species and subspecies of slider turtles, genus *Trachemys*’, *Journal of Herpetology*, vol. 36(2), pp.285–292.

Seidel, ME, Stuart, JN & Degenhardt, WG 1999, 'Variation and species status of slider turtles (Emydidae: *Trachemys*) in the southwestern United States and adjacent Mexico', *Herpetologica*, vol. 55(4), pp. 470–487.

Shepherd, CR & Nijman, V 2007, *An overview of the regulation of the freshwater turtle and tortoise pet trade in Jakarta, Indonesia*, TRAFFIC South East Asia, viewed 22 October 2010, <www.traffic.org/species-reports/traffic_species_reptiles12.pdf>.

Somma, LA, Foster, A & Fuller, P 2009a, *Trachemys scripta elegans*, United States Geological Survey Nonindigenous Aquatic Species Database, viewed 14 October 2010, <<http://nas.er.usgs.gov/queries/FactSheet.aspx?speciesID=1261>>.

Somma, LA, Foster, A & Fuller, P 2009b, *Trachemys scripta scripta*, United States Geological Survey Nonindigenous Aquatic Species Database, viewed 14 October 2010, <<http://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=1262>>.

Spinks, PQ, Pauly, GB, Crayon, JJ & Shaffer HB 2003, 'Survival of the western pond turtle (*Emys marmorata*) in an urban Californian environment', *Biological Conservation*, vol. 113, pp. 257–265.

Stuart, JN 2000, 'Additional notes on native and non-native turtles of the Rio Grande Drainage Basin, New Mexico', *Bulletin of the Chicago Herpetological Society*, vol. 35(10), pp. 229–235.

Telecky, TM 2001, 'United States import and export of live turtles and tortoises', *Turtle and Tortoise Newsletter*, vol. 4, pp. 8–13, viewed 3 November 2010, <<http://www.chelonian.org/ttn/archives/ttn4/pp8-13.shtml>>.

Thomas, AD, Forbes-Faulkner, JC, Speare, R & Murray C 2001, 'Salmonellosis in wildlife from Queensland', *Journal of Wildlife Diseases*, vol. 37(2), pp. 229–238.

TRAFFIC 1998, *The US role in the international live reptile trade: Amazon tree boas to Zululand dwarf chameleons*, TRAFFIC, viewed 22 October 2010, <http://www.traffic.org/species-reports/traffic_species_reptiles3.pdf>.

TRAFFIC 2008, *The state of wildlife trade in China: information on the trade in wild animals and plants in China 2008*, TRAFFIC, viewed 22 October 2010, <www.traffic.org/general-reports/traffic_pub_gen34.pdf>.

Tucker, JK & Paukstis, GL 2000, 'Hatching success of turtle eggs exposed to dry incubation environment', *Journal of Herpetology*, vol. 34, pp. 529–534.

USA Today 2010, *California bans import of turtles and frogs for food*, March 4 2010, USA Today, viewed 22 October 2010, <<http://content.usatoday.com/communities/ondeadline/post/2010/03/calif-bans-imports-of-turtles-frogs-for-food/1>>.

Valdeón, A, Crespo-Díaz, A, Egaña-Callejo, A & Gosá, A 2010, 'Update of the pond slider *Trachemys scripta* (Schoepff, 1792) records in Navarre (Northern Spain), and presentation of the Aranzadi Turtle Trap for its population control', *Aquatic Invasions*, vol. 5(3), pp. 297–302.

Victorian Government 2010, *Weeds and vertebrate pests, module 1: within the invasive plants and animals policy framework*, Department of Primary Industries, Melbourne.

- Vietnam.com 2010, *Clearing up Vietnam's waters from harmful turtles*, 16 October 2010, Vietnam.com, viewed 22 October 2010, <<http://www.vietnam.com/article/clearing-up-vietnams-waters-from-harmful-turtles.html>>.
- Warwick, C 1991, 'Conservation of red-eared terrapins *Trachemys scripta elegans*: threats from international pet and culinary markets', *Testudo*, vol. 3, pp. 34–44.
- Warwick, C, Steedman, C & Holford, T 1990, 'Ecological implications of the red-eared turtle trade', *Texas Journal of Science*, vol. 42(4), pp. 419–422.
- Western Australian Government 2004, *Importing and keeping introduced mammals, birds, reptiles and amphibians in Western Australia*, bulletin 4604, Department of Agriculture, Perth.
- Wikipedia 2010a, *Cumberland slider*, Wikipedia, viewed 26 October 2010, <http://en.wikipedia.org/wiki/Cumberland_slider>.
- Wikipedia 2010b, *Red-eared slider*, Wikipedia, viewed 26 October 2010, <http://en.wikipedia.org/wiki/Red-eared_slider>.
- Williams, T 1999, 'The terrible turtle trade', *Audubon*, vol. 101(2), pp. 44, 46–51.
- World Wildlife Fund (WWF) 2010, *Wildlife trade: reptile trade FAQs*, WWF, viewed 22 October 2010, <<http://www.worldwildlife.org/what/globalmarkets/wildlifetrade/faqs-reptile.html>>.
- Zipcode Zoo 2010, *Trachemys scripta elegans (red-eared terrapin)*, Zipcode Zoo, viewed 21 October 2010, <http://zipcodezoo.com/Animals/T/Trachemys_scripta_elegans/>.