

Blue agave

Agave tequilana



Steve Csurhes
First published 2010
Updated 2016



Queensland
Government

© State of Queensland, 2016.

The Queensland Government supports and encourages the dissemination and exchange of its information. The copyright in this publication is licensed under a Creative Commons Attribution 3.0 Australia (CC BY) licence.



You must keep intact the copyright notice and attribute the State of Queensland as the source of the publication.

Note: Some content in this publication may have different licence terms as indicated.

For more information on this licence visit <http://creativecommons.org/licenses/by/3.0/au/deed.en>

Photo: Wikimedia Commons—licensed under the Creative Commons Attribution 2.0 Generic license).

Contents

Summary	4
Identity and taxonomy	4
Description	5
Native range and global distribution	5
Preferred habitat and climate	5
Reproduction and dispersal	5
History as a weed overseas	5
Use	6
Current distribution and impact in Queensland	6
Potential distribution and impact in Queensland	6
References	6

Summary

A. tequilana (blue agave) is native to the arid highlands of Mexico, where it is cultivated to produce tequila. More recently, there has been some interest in the potential to produce ethanol (“biofuel”) from this plant.

This study was unable to find evidence that *A. tequilana* was a weed anywhere in the world. However, at least 10 congeners have been listed as weeds elsewhere, with three species listed as weeds in southeast Queensland: *A. americana* (century plant), *A. vivipara* and *A. sisalana* (sisal hemp, sisal).

Suitable habitats in Queensland appear to be limited to sandy soils in arid and semi-arid subtropical areas.

Based on the evidence collected in this preliminary study, *Agave tequilana* appears to pose a low weed risk.

Identity and taxonomy

Species: *Agave tequilana* FAC Weber (includes commercial var. “azul”)

Other synonyms:

Agave palmaris Trel.

Agave pedrosana Trel.

Agave pes-mulae Trel.

Agave pseudotequilana Trel.

Agave subtilise Trel.

Common names:

“blue agave”, “Weber’s blue agave” or “tequila weed”

Family: Agavaceae

Description

Like other *Agave* species, *A. tequilana* grows as a large rosette of thick, fleshy leaves, each ending in a sharp point (see cover photo).

Native range and global distribution

A. tequilana is native to the central western highlands of Mexico (Jalisco, Oaxaca and Puebla) (USDA 2009).

Preferred habitat and climate

Within its native range, *A. tequilana* favours sandy soils in arid or semi-arid areas. While Mexico and Central America lie in the tropics, *A. tequilana* grows at altitudes of more than 1,500 metres (m), where climate is considerably cooler than lowland areas.

Reproduction and dispersal

Each plant is monocarpic, flowering only once (at about five years old). During flowering, a “mast” (stem), up to five metres tall, grows from the centre of the plant and bears a large number of tubular, yellow flowers.

Within its native range, *A. tequilana* is pollinated by a locally native bat (*Leptonycteris nivalis*) and produce several thousand seeds per plant. The plant then dies.

Agave species do not appear to produce seeds in Australia, perhaps due to an absence of suitable pollinators. Reproduction of *Agave* species in Australia is believed to be restricted to vegetative reproduction.

Specimens have been recorded to live for 50 years in gardens.

History as a weed overseas

A. tequilana is listed as “introduced” by Randall (2002), but not naturalised or weedy. However, Randall listed 10 congeners: *A. americana*, *A. angustifolia*, *A. attenuata*, *A. decipiens*, *A. desmettiana*, *A. lecheguilla*, *A. schottii*, *A. sisalana*, *A. vivipara* and *A. weberi* as either naturalised or weedy.

Holm et al. (1979) listed *A. lecheguilla* and *A. sisalana* as weeds.

A. americana (century plant), *A. vivipara* and *A. sisalana* (sisal hemp, sisal) are naturalised in southeast Queensland (Queensland Herbarium 2007). Perhaps the most problematic species are *A. americana* and *A. sisalana*, which form small clumps close to where they were originally planted. Such clumps are sometimes impenetrable due to their strong thorns. Control is sometimes undertaken within conservation reserves (eg. some island national parks in north Queensland).

Use

A. tequilana var. “azul” is the base ingredient of tequila, a popular alcoholic drink, and an important economic product of Mexico where some 800,000 ton is processed annually (Ballinas et al. 2008). Some 200 million specimens are cultivated in the western central highlands of Mexico.

Tequila is produced by removing the heart of the plant in its twelfth year, normally weighing between 35–90 kg. This heart is stripped of leaves and heated to remove the sap, which is fermented and distilled. Other beverages like mezcal and pulque are also produced from blue and other agaves by different methods (though still using the sap) and are regarded as more traditional.

More recently, there has been some interest in the potential to grow *A. tequilana* as a source of biofuel (ethanol).

Current distribution and impact in Queensland

A. tequilana is not listed as naturalised in Queensland (Queensland Herbarium 2007). However, it is cultivated as an ornamental.

Potential distribution and impact in Queensland

This study was unable to find evidence that *A. tequilana* was a weed anywhere in the world. However, at least 10 congeners have been listed as weeds elsewhere, with three species listed as weeds in southeast Queensland. The latter species are generally considered to be minor weeds, despite being capable of forming dense, localised thickets (spread is slow, due probably to the absence of viable seeds).

A. tequilana is perhaps best suited to sandy soils in arid and semi-arid subtropical areas.

References

Ballinas, L, Manjarrez, L, Fierro, V, Goncalves, A, Moraes-Rocha, G and Gonzalez-Sanchez, G (2008). Microscopic characterisation of *Agave tequilana weber* var. azul fibers from agroindustrial waste in activated carbon production. *Microscopy and Microanalysis* 14 (Suppl 2). Microscopy Society of America.

Holm, LG, Pancho, JV, Herberger, JP. and Plucknett, DL (1979). ‘A Geographical Atlas of World Weeds’ (Krieger Publishing Company, Florida).

Queensland Herbarium (2007). Census of the Queensland Flora. Edited by P Bostock and A. Holland, Queensland Government, Brisbane.

Randall, RP (2002). A Global Compendium of Weeds, RG and FJ Richardson, Melbourne.

USDA (2009). ARS, National Genetic Resources Program, Germplasm Resources Information Network–(GRIN) [Online Database], National Germplasm Resources Laboratory, Beltsville, Maryland. URL: <http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?1738> (10 February 2009).